

# Ankle Arthrodesis

OS14 Lite - Expires end of January 2022



This fact sheet is for general information about this procedure only. It is not intended to be used as medical advice or to replace advice that your relevant healthcare professional would give you. If you have a particular medical problem, please consult a healthcare professional.

© EIDO Systems International Limited. The operation and treatment information in this document is published under license from EIDO Systems International and is protected by copyright laws. Other than for your personal, non-commercial use, you may not copy, print out, download or otherwise reproduce any of the information.

[eidohealthcare.com](http://eidohealthcare.com)



## What is arthritis?

Arthritis is a group of conditions that cause damage to one or more joints.

The most common type of arthritis is osteoarthritis, where there is gradual wear and tear of a joint.

Some other types of arthritis are associated with inflammation of the joints that can eventually lead to severe joint damage.

Arthritis eventually wears away the normal cartilage covering the surface of the joint and the bone underneath becomes damaged.


## What are the benefits of surgery?

You should get less pain and be able to walk more easily.

## Are there any alternatives to ankle arthrodesis?

Simple painkillers such as paracetamol and anti-inflammatory painkillers such as ibuprofen can help control the pain of arthritis.

Using a walking stick on the opposite side to the affected ankle can make walking easier. A plastic splint or a stiff ankle boot with a cushioned heel is sometimes helpful. Physiotherapy may help to strengthen weak muscles.



An arthritic ankle

A steroid injection into your ankle joint can sometimes reduce pain and stiffness for several months.

An arthroscopy (keyhole surgery) to clean out your ankle joint can give some relief for 6 to 12 months.

All these measures become less effective if your arthritis gets worse and this is when your surgeon may recommend an ankle arthrodesis.

Some people with ankle arthritis can have an ankle replacement instead of an ankle arthrodesis.

## What does the operation involve?

Various anaesthetic techniques are possible. The operation usually takes an hour to 90 minutes.

Your surgeon will remove the damaged joint surfaces. They will fix the bones together with a metal plate, a metal rod, or screws.

## What complications can happen?

Some complications can be serious and can even cause death.

### General complications of any operation

- Pain
- Bleeding
- Infection of the surgical site (wound)
- Allergic reaction to the equipment, materials or medication
- Unsightly scarring of your skin
- Blood clot in your leg
- Blood clot in your lung
- Difficulty passing urine

### Specific complications of this operation

- Damage to nerves around your ankle
- Breakdown of the skin over your ankle
- Infection in your ankle
- Failure of the arthrodesis, where the bones do not join together
- Severe pain, stiffness and loss of use of your foot and ankle

## How soon will I recover?

You should be able to go home after 1 to 2 days.

To start with, spend most of the time with your leg raised on a chair or footstool.

Most people need to have the cast for about 8 to 12 weeks. You will need to use a walking aid until it is removed. Once the cast is removed, regular exercise should help you to return to normal activities as soon as possible. Before you start exercising, ask the healthcare team or your GP for advice.

Most people make a good recovery, have less pain, and can move about better.

## Summary

If you have severe arthritis in your ankle, an ankle arthrodesis should reduce your pain and allow you to do more of your normal activities.

Keep this information document. Use it to help you if you need to talk to the healthcare team.

Some information, such as risk and complication statistics, is taken from global studies and/or databases. Please ask your surgeon or doctor for more information about the risks that are specific to you.

This document is intended for information purposes only and should not replace advice that your relevant healthcare team would give you.

### Acknowledgements

Reviewer: Stephen Milner (DM, FRCS (Tr & Orth))

Illustrator: Medical Illustration Copyright ©

Medical-Artist.com